

Effect of Intrathecal Dexmedetomidine and Fentanyl as Adjuvant to Hyperbaric Bupivacaine for Orthopaedic Lower Limb and Lower Abdominal Procedures: A Double Blind Control Study

Sidharth Srabam Routray^{1,*}, Khageswar Ravi², Debasis Mishra³

^{1,2,3}Assistant Professor, Dept. of Anaesthesiology, SCB Medical College, Cuttack, Odisha

*Corresponding Author:

E-mail: drkitusraban@gmail.com

ABSTRACT

Aims and Objectives: The aim of this study is to evaluate the effects of Dexmedetomidine and Fentanyl as intrathecal adjuvant to Bupivacaine on the onset and duration of sensory and motor block in spinal anaesthesia in orthopaedic lower limb and lower abdominal surgeries.

Materials and Method: Ninety patients of ASA status I and II posted for lower limb and lower abdominal surgery were randomly divided into three groups. Group D was administered Hyperbaric Bupivacaine 15mg + Dexmedetomidine 5µg in 0.5 ml normal saline, group F was administered Hyperbaric Bupivacaine 15mg + Fentanyl 25 µg in 0.5 ml normal saline and group C were administered Hyperbaric Bupivacaine 15mg plus in 0.5 ml normal saline. Duration and quality of sensory and motor block were assessed.

Results: Sensory and motor block in group D patients were longer than group F and C patients.

Conclusion: Intrathecal dexmedetomidine when added to bupivacaine heavy (0.5%) provide better and prolonged analgesia in comparison to fentanyl.

Key word: Fentanyl, Orthopaedic, Bupivacaine, Intrathecal, Dexmedetomidine

Access this article online	
Quick Response Code:	Website: www.innovativepublication.com
	DOI: 10.5958/2394-4994.2015.00030.X

INTRODUCTION

Subarachnoid block is the most commonly used technique for orthopaedic lower limb and lower abdominal surgeries^[1]. Postoperative pain control is a major problem in these surgeries because of relatively short duration of action of local anaesthetics, so early analgesic intervention is needed in the postoperative period. A number of adjuvants, such as clonidine and midazolam, and opioids have been studied to prolong the effect of spinal anaesthesia^[2,3]. Fentanyl, in recent years, has emerged as a useful intrathecal adjuvant for prolonging the effect of spinal anaesthesia. Although it is one of the most widely used intrathecal adjuvant in the present scenario, its intrathecal use has been shown to be associated with side effects like respiratory depression and pruritus^[4].

Dexmedetomidine, a new highly selective α -2-agonist, is under evaluation as a neuraxial adjuvant as it provides stable hemodynamic conditions, good quality of intraoperative and prolonged postoperative analgesia with minimal side effects. By virtue of its effect on spinal α -2 receptors, dexmedetomidine mediates its

analgesic effects. Based on earlier human studies, it has been shown that a low dose of 5 µg, dexmedetomidine provides a prolonged anaesthesia and good quality post operative analgesia when used as an intrathecal adjuvant to bupivacaine with minimal effects on the hemodynamic status of the patient^[5,6].

Hence, the present study is being undertaken to evaluate and compare the effects of dexmedetomidine and fentanyl as intrathecal adjuvants to bupivacaine.

MATERIAL AND METHODS

This study was carried out at SCB Medical College & Hospital, Cuttack after obtaining approval from the Hospital Ethical Committee and written informed consent from the patients from July 2014 to Feb. 2015. Ninety ASA grade I or II patients of either sex, aged 18 to 60 years, weighing 50 to 90 kg and with a height of 150 cm to 180 cm, scheduled for lower abdominal and lower limb surgery were included in the study. Exclusion criteria were Patients presenting with known contra indications to spinal anaesthesia, pregnant patients, Patients on therapy with adrenergic receptor antagonist, calcium channel blocker, and/or ACE inhibitor, with history of heart block or dysarrhythmia, hypersensitivity to any of the study drugs, who refused to consent to be part of study.

The study population was randomized using random number table generated from computer software. Random intervention assignment slip was placed in serially numbered opaque and sealed

envelopes. These envelopes were opened following enrolment of the case.

90 total patients randomly divided into 3 groups (n = 30): Group D: Hyperbaric Bupivacaine 15mg + Dexmedetomidine 5µg in 0.5 ml normal saline administered intrathecally. Group F: Hyperbaric Bupivacaine 15mg + Fentanyl 25 µg in 0.5 ml normal saline administered intrathecally. Group B: Hyperbaric Bupivacaine 15mg plus 0.5 ml normal saline administered intrathecally.

All the patients were kept for 6 hrs fasting prior to surgery. Tablet Alprazolam (0.25 mg) was given as a premedication a night prior to surgery. Preloading was done with Ringer lactate solution (10 ml/kg body weight). Routine monitoring including non invasive blood pressure (NIBP), ECG, heart rate and pulse oximetry was done. All patients received supplemental oxygen via mask (3l/min).

Under proper aseptic conditions, spinal anaesthesia was given at the level of L3-L4 interspace in sitting position using a midline approach by a 25G Quincke spinal needle. The drug was injected slowly over 10-15 seconds with the bevel of the needle pointing upwards and all patients were made supine.

The intrathecal drug formula was prepared by a separate anaesthesiologist under strict aseptic conditions. The anaesthesiologist who administered anaesthesia was blinded to the group allocation. After administering anaesthesia the vital signs of the patient were recorded. Vitals were recorded every 2 minutes up to the 10th minute and every 5 minutes thereafter up to 20 minutes. Beyond 20 minutes the vitals were recorded every 20 minutes till the time of discharge from PACU (Post Anaesthesia Care Unit). The sensory dermatome level was assessed by loss of pin prick sensation to a 23 G hypodermic needle.

The motor dermatome level was assessed according to the Bromage^[7] Scale:

- Bromage 0- Patient able to move hip, knee and ankle.
- Bromage 1- Patient unable to move hip, but able to move knee and ankle.
- Bromage 2- Patient unable to move hip and knee but able to move the ankle.
- Bromage 3- Patient unable to move hip, knee and ankle.

The sensory and motor status was assessed prior to the spinal injection, then every 2 minutes after the spinal injection for the first 10 minutes, every 5 minutes for the next 10 minutes and thereafter every 20 minutes until the time to regression of sensory level to dermatome S2 and motor scale to bromage 0.

Time to reach the sensory block up to highest dermatome level and motor block of bromage 3 level was noted. On achieving T8 sensory blocked level, the surgical procedure was carried out. Then time to

regression to dermatome S2 level and time to reach bromage 0 was noted in post operative care unit. All durations were calculated taking the spinal injection time as time zero. If the sensory levels were not equal bilaterally the higher dermatome level was used for statistical analysis. Patient was discharged after the sensory block regresses to S2 level and motor block to bromage 0.

Postoperatively, the pain scoring was done by using visual analog scale^[8] (VAS) (0 = no pain, 10 = severe pain), with the vital recordings of the study until the patient was discharged. Paracetamol was given intravenous as rescue analgesia when VAS was greater than 4. Time of administering the first dose of rescue analgesia was noted. Sedation was assessed by using Modified Ramsay sedation score each time the vitals were noted.

Modified Ramsay sedation scale^[9]:

1. Anxious, Agitated, Restless.
2. Cooperative, Oriented, Tranquil.
3. Responds to commands only.
4. Brisk response to light glabellar tap or loud noise.
5. Sluggish response to light glabellar tap or loud noise.
6. No Response.

For the purpose of the study hypotension was defined as a decrease in systolic blood pressure more than 30% of the baseline value or fall below 90 mmHg, which was treated by Ephedrine 6 mg i.v. and fluids. Bradycardia was defined as heart rate less than 60/min but the intervention with iv atropine 0.6mg was done only when heart rate fell below 50/min.

Side effects including nausea, vomiting, bradycardia, hypotension, pruritus, respiratory depression, urinary retention, shivering etc. were assessed both intra-operatively as well as post-operatively. All the patients were examined by the anaesthesiologist after 24 hrs of the spinal block and were assessed for any postdural puncture headache or transient neurologic symptoms. Highest dermatomal level of sensory blockade, time taken to reach the highest dermatomal level of sensory block, to reach up to bromage 3 motor block, for sensory regression to S2 level, for motor regression to bromage 0 were noted. Hemodynamic status of the patient, sedation score and side effects if any were noted.

All the statistical analysis was performed by using SPSS version 21. The various statistical tests that were used in this study were analysis of variance (ANOVA) test, Post hoc test (Bonferroni test) and nonparametric tests like Mann whitney U test and kruskalwallis test. For all statistical analysis p < 0.05 was considered statistically significant. The results and interpretations are explained below.

RESULT

All the three groups were comparable as regard to age, sex, height and weight. There were no significant differences in heart rate, MAP, SPO₂ between the three groups. Intergroup analysis showed a statistically significant difference in the highest level of sensory blockade amongst group D and F ($p = 0.004$). However no significant difference was found between group B and F and group D. Two segment regression time was more in dexmede-tomidine group in comparison to fentanyl and control group.

Onset time to both sensory and motor block

was faster on group D than group F and group B. Regression time of motor block to bromage 0 was slow and time to rescue analgesia was longer in dexmedetomidine group in comparison to other groups. Sedation score was more in dexmede-tomidine group in comparison to other group. VAS score was lower in dexmedetomidine group than other groups. In our study the incidence of bradycardia, hypotension, nausea, vomiting, pruritus and urine retention was not statistically significant. There was no incidence of respiratory depression.

Table 1: Demographic profile

Parameters	Group B	Group D	Group F	P value
Age (years)	39.16±10.12	43.6±10.5	41.63±9.85	0.245
Sex(m:f)	20:10	19:11	17:13	0.548
Weight (kg)	66.87±7.20	66±7.82	64.83±7.50	0.577
Height (cm)	166.47±6.40	164.43±5.92	163.6±5.62	0.168
ASA(I:II)	21:9	18:12	20:10	0.641
Duration of surgery (min)	48±17.84	47.67±15.24	46.47±11.99	0.734

Table 2: Characteristics of spinal block

Variable (min)	Group B	Group D	Group F	P value
Time of onset of sensory block	8.2±1.6	8.0±1.5	8.3±2.1	0.235
Time of onset of motor block	9.5±2.3	9.3±2.7	9.7±3.2	0.124
Time taken to reach highest level of sensory block	10.4±4.01	9.33±3.50	10.67±3.65	0.346
Duration of sensory block	198.67±32.35	396.67±24.12	205.67±26.12	0.0001
Duration of motor block	140.67±21.32	338±21.24	154±19.76	0.0001
Duration of Spinal anesthesia	224.4±36.2	402.6±32.7	234.6±29.6	0.0001
Time to first dose of rescue analgesia	153.67±27.88	299±33.92	166.83±20.66	0.0001

Table 3: Highest dermatome level of sensory block

HIGHEST LEVEL OF SENSORY DERMATOME	GROUPS			TOTAL
	B	D	F	
T4	3	4	0	7
T5	9	14	6	29
T6	8	5	11	24
T7	2	4	6	12
T8	5	2	3	10
T9	3	1	3	7
T10	0	0	1	1

	Mann Whitney U test		Kruskalwallis (p value)
B & D	B & F	D & F	
0.149	0.191	0.004	0.017

Table 4: Adverse effect of spinal block, (values are numbers)

Side effect	Group B	Group D	Group F	P value
Hypotension	4	3	3	0.895
Bradycardia	2	4	3	0.693
Resp. depression	0	0	0	1
Nausea, vomiting	3	1	2	0.589
Urinary retention	1	3	1	0.433
Pruritus	1	0	2	0.36
Dry mouth	3	2	2	0.858
Shivering	4	1	2	0.342
Sedation score	1.35±0.07	1.50±0.09	1.44±0.08	0.0001

Table 5: Postoperative visual analogue scale (VAS)

Time after surgery	Group B	Group D	Group F
1 hr	4(2-5)	0(0-3)	4(2-4)
2hr	4(1-4)	1(0-3)	3(2-4)
3hr	3(1-4)	1(0-2)	3(1-3)
4hr	3(1-3)	2(0-3)	2(0-3)
5hr	2(1-3)	2(0-2)	2(0-3)

DISCUSSION

The results of our study show that the supplementation of intrathecal bupivacaine with 5µg dexmedetomidine significantly prolonged both sensory and motor block compared with intrathecal 25 µgfentanyl and control group. Patients in the groups that received dexmedetomidine and fentanyl had reduced postoperative pain scores and a longer pain free period than those who received spinal bupivacaine alone. No hemodynamic instability or adverse effects were reported in any group. Time taken to achieve peak level of sensory and motor blockade was earlier in dexmedetomidine group than other groups.

Al Ghanem et al^[10] observed that the onset time of bromage 3 motor block was also not different between dexmedetomidine and fentanyl group. Regarding time taken to achieve peak motor blockade there was no statistically significant difference was seen amongst all the three groups. The time to regression of sensory block to S1 segment was significantly longer in group D than in group F ($p < 0.001$). The regression time to reach bromage 0 in dexmedetomidine group was significantly longer than that for fentanyl group ($p < 0.001$). This was similar to our study in which dexmedetomidine group showed a statistically significant prolongation of both sensory and motor regression when compared to fentanyl and bupivacaine alone group.

Al-Mustafa M M et al^[11] studied the effect of adding different doses of dexmedetomidine (5µg or 10µg) to bupivacaine (12.5mg) for neuraxial anaesthesia. They observed a maximum sedation score of 2 without pre-medicating their patients with any type of benzodiazepines in both the groups. In our study the mean blood pressure in the post operative period, was

found to be similar all groups. Kanazi et al^[12] noted that dexmedetomidine or clonidine when added to intrathecal bupivacaine did not cause a significant reduction in blood pressure but prolonged sensory and motor block characteristics.

El-Hennawy AM et al^[13] found that addition of clonidine or dexmedetomidine to bupivacaine prolongs caudal analgesia in children. Alka Shah et al^[14] studied Haemodynamic effects of intrathecal dexmedetomidine added to ropivacaine intraoperatively and found that it prolonged the postoperative analgesia. Gehan A. Tarbeehet al^[15] studied the effects of intrathecal bupivacaine-fentanyl versus bupivacaine-dexmedetomidine in diabetic surgical patients and concluded that dexmedetomidine produced better block characteristics. Jain et al^[16] studied the perioperative effect of epidural dexmedetomidine with intrathecal bupivacaine on haemodynamic parameter and quality of analgesia and found that it is better than other adjuvants.

Bradycardia was seen in total of 9 patients in our study and was more in dexmedetomidine group (4 patients) compared to fentanyl and bupivacaine alone group, but it was transient and did not require any intervention. There was no statistically significant difference noted amongst the three groups. Urinary retention was seen in 3 patients in dexmedetomidine group compared to one patient each in the other two groups but it was statistically not significant ($p > 0.05$).

Nausea and vomiting was highest in bupivacaine alone group followed by fentanyl group and least in dexmedetomidine group. It was also not statistically significant on analysis ($p > 0.05$). The administration of intrathecal opioids may provide benefits in augmenting intra operative anaesthesia but

carries a risk of respiratory depression. Varassi et al^[17] demonstrated that intrathecal administration of fentanyl 25 micrograms in non premedicated geriatric patients did not alter respiratory rate, ETCO₂, minute ventilation, respiratory drive and ventilator response to CO₂. On the contrary, 50µg intrathecal fentanyl can cause an early respiratory depression in geriatric patients. In our study none of the groups showed any effect on respiratory rate or any decrease in O₂ saturation.

CONCLUSION

We conclude from our study that supplementation of bupivacaine spinal block with a low dose of 5µg intrathecal dexmedetomidine produces a significantly longer duration of sensory and motor block than 25µg intrathecal fentanyl. It provides hemodynamically stable conditions, minimal side effects, and excellent quality of postoperative analgesia. Thus, 5 µg dexmedetomidine seems to be an attractive alternative to 25 µg fentanyl as an adjuvant to spinal bupivacaine in surgical procedures

REFERENCES:

- Mulroy M F. Regional anaesthetic techniques. *IntAnesthesiolClin* 1994; 32(3):81-98.
- Elia N, Culebras X, Mazza C, Schiffer E, Tramèr MR. Clonidine as an adjuvant to intrathecal local anesthetics for surgery: Systematic review of randomized trials. *RegAnesth Pain Med* 2008;33:159-67.
- Bousofara M, Carlès M, Raucoules-Aimé M, Sellam MR, Horn JL. Effects of intrathecal midazolam on postoperative analgesia when added to a bupivacaine-clonidine mixture. *RegAnesth Pain Med* 2006;31:501-5.
- Shende D, Cooper GM, Bowden MI. The influence of intrathecal fentanyl on the characteristics of subarachnoid block for caesarean section. *Anaesthesia* 1998; 79: 702-710.
- Anand VG, Kannan M, Thavamani A, Bridgit MJ. Effects of dexmedetomidine added to caudal ropivacaine in paediatric lower abdominal surgeries. *Indian J Anaesth* 2011;55:340-6
- Coursin D B, Maccioli G A. Dexmedetomidine. *CurrOpinCrit Care* 2001; 7: 221-6
- Robert K Stoelting. Opioid agonists and antagonists. In *Pharmacology and Physiology in anesthetic practice*. Ed by Robert K Stoelting, Simon C Hillier. 4th edn. Lippincott Williams and Wilkins. 1999; 87-122.
- Panzer O, Moitra V, Roberet N Sladen. Pharmacology of sedative-analgesic agents – Dexmedetomidine, remifentanyl, ketamine, volatile anaesthetics and the role of Mu antagonists. *Critical Clin* 2009;25:451-69.
- Ralph Getler, Clieghton H Brown, Mitchel H, Silvius N. Dexmedetomidine: a novel sedative analgesic agent. *Baylor University Medical Centre Proceedings*. 2001;14(1).
- Al-Ghanem SM, Massad IM, Al-Mustafa MM, Al-Zaben KR, Qudaisat IY, Qatawneh AM and Abu-Ali HM. Effect of Adding Dexmedetomidine versus Fentanyl to Intrathecal Bupivacaine on Spinal Block Characteristics in Gynecological Procedures: A Double Blind Controlled Study. *Am J ApplSci* 2009;6:882-7.
- Al-Mustafa MM, Abu-Halaweh SA, Aloweidi AS, Murshidi MM, Ammari BA, Awwad ZM, et al. Effect of dexmedetomidine added to spinal bupivacaine for urological procedure. *Saudi Med J* 2009;30:365-70.
- Kanazi GE, Aouad MT, Jabbour-Khoury SI, Al Jazzar MD, Alameddine MM, Al-Yaman R, et al. Effect of low-dose dexmedetomidine or clonidine on the characteristics of bupivacaine spinal block. *ActaAnesthesiolScand* 2006;50:222-7.
- El-Hennawy AM, Abd-Elwahab. Addition of clonidine or dexmedetomidine to bupivacaine prolongs caudal analgesia in children. *Br J Anaesth* 2009;103:268-74.
- Alka Shah, Ila Patel, Rachana Gandhi et al. Haemodynamic effects of intrathecal dexmedetomidine added to ropivacaine intraoperatively and for postoperative analgesia. *International Journal of Basic & Clinical Pharmacology* | January-February 2013 | Vol 2 | Issue 1, 26-29.
- Gehan A, Tarbeeh, Ayman A, Mohamed et al. Effects of intrathecal bupivacaine-fentanyl versus bupivacaine-dexmedetomidine in diabetic surgical patients. *Egyptian journal of Anaesthesia*(2013) 29, 13-18.
- Jain D, Kumar D, Khan RM, Kumar N et al. Perioperative effect of epidural dexmedetomidine with intrathecal bupivacaine on haemodynamic parameter and quality of analgesia. *South Afr J Anaesth Analg* 2012;18(1):105-109.
- G.Varrassi, D.Celleno. G.Caponga et al. "Ventilatory effects of subarachnoid Fentanyl in elderly", *Anaesthesia* 1992, Vol47: 558-562.