

Comparison of Caudal Bupivacaine with Additives Fentanyl or Ketamine for Post-Operative Pain Relief in Children: A Randomized Controlled, Double Blinded Study

Sawai Singh Jaitawat^{1,*}, Rakesh Kushwaha², Nagina³, Sunanda Gupta⁴

¹Professor, ²Associate Professor, ³Resident, ⁴Professor & HOD, Department of Anaesthesiology and Critical Care, Geetanjali Medical College & Hospital Udaipur, Rajasthan, India

*Corresponding Author:

E-mail: Jaitawatss@gmail.com

ABSTRACT

Background and Aims: Ensuring adequate analgesia in the post-operative period is an indispensable part of a balanced anesthesia technique with increasing scope of day care surgery and emphasis on early discharge. In children undergoing infra-umbilical and lower limb surgeries, caudal block is a reliable and safe technique that can be used with general anesthesia for intra and post-operative analgesia. The present study was designed to compare the analgesic efficacy of ketamine and fentanyl as additives to bupivacaine given caudally in children undergoing lower abdominal and lower limb surgery.

Materials and Methods: A total of 60 children aged 1-5 years undergoing lower abdominal and lower limb surgery were included in this randomized, controlled, double-blind study. Three groups of 20 each were assigned to receive caudal block with bupivacaine 0.25% 0.5 ml/kg alone (group A) or along with 1 ml/kg fentanyl (group B) or 0.5 mg/kg ketamine (group C). Assessment of post-operative pain was done using the Objective Pain Scale. Requirement of rescue analgesia and side effects were also noted.

Results: Children who received ketamine with bupivacaine caudally (group C) had the longest duration of post-operative analgesia [Group A vs. Group C: 150 min ($p=0.32$) vs. 180 min ($p=0.00$)] and requirement of first dose of analgesia in groups A, B and C ranged from 183 ± 16.5 min, 350 ± 2.5 min and 594 ± 148.12 min respectively. The mean number of doses of rescue analgesia that patients received in 24 hours in group A, B and C were 6.0, 1.63 and 1.40 respectively.

Conclusion: Caudally administered Ketamine in the dose of 0.5 mg/kg with bupivacaine provides prolonged post-operative analgesia in comparison to fentanyl 1 μ g/kg with bupivacaine with minimal side effects.

Keywords: Ketamine, fentanyl, bupivacaine, children

Access this article online	
Quick Response Code:	Website: www.innovativepublication.com
	DOI: 10.5958/2394-4994.2015.00038.4

INTRODUCTION

Ensuring adequate analgesia in the post-operative period is an indispensable part of a balanced anesthesia technique with increasing scope of day care surgery and emphasis on early discharge. Patients can experience pain after discharge when suitable medical help is unavailable. In children undergoing infra-umbilical and lower limb surgeries, caudal block is a reliable and safe technique that can be used with general anesthesia for intra and post-operative analgesia⁽²⁾. It is the one of most commonly used regional technique for post-operative analgesia in children⁽³⁾. Post-operative pain relief is commonly provided by parenteral or oral drugs including opioids and non-steroidal anti-inflammatory drugs (NSAIDs). Inability to ensure adequate pain relief due to variability and individual sensitivity is a considerable drawback

of this approach. Some of these drugs such as oral NSAIDs have been reported to have side effects like gastrointestinal hemorrhage and renal papillary necrosis.⁽⁴⁾ Therapeutic doses of opioids are not administered due to concerns of respiratory depression, nausea, vomiting, and distressing pruritus.⁽⁵⁾ Many anesthetic agents have been used for caudal analgesia in pediatric patients. Bupivacaine and Levobupivacaine have proven clinical effectiveness and safety.⁽⁶⁻⁸⁾ A number of adjuvants such as fentanyl, clonidine, Ketamine and, midazolam have been added to local anesthetic solutions to prolong caudal analgesia as a single bolus⁽⁹⁻¹⁴⁾. A combination of ketamine and bupivacaine increases the potency and duration of analgesia⁽¹⁵⁻¹⁶⁾. Caudal fentanyl in a dose of 1 mcg/kg in addition to bupivacaine has been associated with decreased heart rate due to the systemic effect of fentanyl⁽¹⁷⁾. The present study aims to compare the hemodynamic and analgesic effects of caudal bupivacaine along with fentanyl and ketamine as adjuvants in children aged 1 to 5 years undergoing lower limb or genitourinary surgery.

MATERIALS AND METHODS

This prospective double-blinded study was conducted in the department of anaesthesiology and

critical care of Geetanjali Medical College, Udaipur, Rajasthan which is a tertiary care hospital. After approval of the research protocol by the institutional ethics committee and informed consent from the legal guardians of 60 ASA status I children between 1-5 years of age undergoing lower limb or genitourinary surgery were randomly divided into three groups of 20 each:

Group A received 0.25% bupivacaine as 0.5 ml/kg. Group B received 0.25% bupivacaine as 0.5 ml/kg with fentanyl 1 mcg/kg. Group C received 0.25% bupivacaine as 0.5 ml/kg with ketamine 0.5 mg/kg. Normal saline was added to make equal volumes of the mixtures.

The following children were excluded from the study:

- a. Bleeding/coagulation disorder
- b. Hepato-renal disease
- c. Cardiovascular disease
- d. Local deformity/infection in sacral region
- e. Neurosurgical spinal disorder
- f. Known allergy to local anesthetics
- g. Body weight >25 kg

All the children were examined a day prior to surgery. Pre-operatively hemoglobin estimation and urine examination for albumin and sugar were performed. All children were scheduled to receive general anesthesia combined with caudal block. In the pre-operative hold area, peripheral intravenous access was secured and all children were administered inj. ketamine 2mg/kg and inj. glycopyrrolate 0.02 mg/kg. To facilitate endotracheal intubation inj. Succinylcholine 1.5 mg/kg was used. For maintenance of anesthesia, mixture of oxygen, nitrous oxide, sevoflurane and intermittent doses of inj. atracurium were used. Intraoperatively, the children received 5% dextrose in 0.33% normal saline at 4 ml/kg/hour.

Caudal anesthesia was administered in the lateral position with a 22G hypodermic needle and one of the three study drugs solution were administered as for random allocation of the groups while the other authors who gave caudal block were blinded to the study groups. Two authors were involved in preparing the above mentioned mixtures. Heart rate, systolic blood pressure (SBP), SpO₂, awareness (sweating, tears) was recorded intraoperatively. The parameters recorded post-operatively were pulse rate, blood pressure, SpO₂. The Objective Pain Scale developed by Hannallah and colleagues⁽¹⁸⁾ was used for the assessment of pain.

At the end of surgery, neuromuscular blockade was reversed using inj. neostigmine 0.05mg/kg and inj. atropine 0.02 mg/kg followed by extubation. The children were monitored for vomiting, pruritus and respiratory depression and discharged from the recovery room when an Aldrete score of 10 was achieved⁽¹⁸⁾. The time of first requirement of rescue

analgesic was recorded. Pain scores were assessed at 15 minutes, 1 hour, 4 hours and 8 hours 12 hours. After surgery. When the patient had a pain score of >6, oral acetaminophen 15-20 mg/kg was administered. Vomiting and time of the first micturition were noted along with the total number of acetaminophen doses of rescue analgesia in 24 hours.

STATISTICAL ANALYSES

The data was analysed applying the student's t-test Chi square test and Analysis of variance (ANOVA) where applicable data was expressed as mean + standard deviation and $P < 0.05$ was considered to be significant.

RESULTS

The patients in all three groups were comparable with respect to age and body weight (Table 1). The surgical procedures that the children underwent were repair of hypospadias, inguinal Herniotomy, orchidopexy, circumcision and correction of congenital dislocation of the hip. On comparison, there was no statistical difference between the groups with respect to the type and duration of surgery and the volume of the study drugs.

Intraoperative Hemodynamic Parameters: There was no significant change in heart rate immediately after surgical incision and Intraoperatively in group A and group C. Group B had a significant decrease in heart rate after incision that persisted till 10, 15 and 20 minutes (p value 0.12, 0.001 and 0.00 respectively). After 20 minutes, there was no significant difference in heart rate in all three groups. No significant change was observed in any of the groups with respect to systolic and diastolic blood pressure, and oxygen saturation. All three groups were comparable in pain scores in the first 3 hours after surgery (Table 2). Group A had significantly higher pain scores as compared to group B at 150 and 180 minutes. Patients in group B had significantly higher pain score in comparison to Group C at 310 and 360 minutes. The mean pain score in group C remained low up to 24 hours.

Rescue analgesia: The duration of post-operative analgesia was significantly higher in group C in comparison to group A and group C. The mean number of doses of rescue analgesia that patients received in 24 hours in group A, B and C were 6.0, 1.63 and 1.40 respectively.

Side effects: One patient in group B had an episode of vomiting in the recovery room 20 minutes after surgery which was treated with intravenous metoclopramide. Two patients in group C had 'vacant stares' up to one hour which was not apparently distressing to the parents and resolved spontaneously in about 2 hours.

Table 1: Demographic Table

	Age in year (Mean+SD)	Weight in kg (Mean+SD)
Group A	3.25+1.20	13.35+ 2.05
Group B	2.97+1.40	12.95+3.98
Group C	3.45+1.27	14.28+3.43
F Value	0.692	0.869
P Value	0.505	0.425

Table 2: Post Operative Objective Pain Score(Hanallah)

Time		15min.	1 hrs.	4hrs.	8hrs	12hrs	24hr.
Group A	Mean	1.15	1.05	5.50	6.00	7.00	7.00
	SD	.933	.826	1.150	.000	.000	.00
	Median	1.00	1.00	6.00	6.00	6.5	6.5
Group B	Mean	.90	1.30	2.25	2.25	6.00	6.00
	SD	.308	.470	.639	.605	.00	.00
	Median	1.00	1.00	2.00	2.00	6.00	6.00
Group C	Mean	1.30	1.35	2.10	2.10	2.55	4.62
	SD	.470	.489	.852	.632	.51	1.68
	Median	1.00	1.00	2.00	2.00	3.00	6.00

P Value: Group A vs. B: 0.00 (150 minute), 0.001 (180 minute)
 Group A vs. C: 0.32 (150 minute), 0.00 (180 minute)
 Group B vs. C: 0.00 (310 minute), 0.00 (360 minute)

Table 3: Objective Pain Scale (OPS) of Hanallah et al for Post-Operative pain Assessment

Parameter	Finding	Points
Systolic blood pressure	Increase<20% of preoperative blood pressure	0
	Increase<20-30% of preoperative blood pressure	1
	Increase>30% of preoperative blood pressure	2
Crying	Not crying	0
	Respond to age appropriate nurturing (tender loving care)	1
	Does not respond to nurturing	2
Movements	No movements relaxed	0
	Restless moving about in bed constantly	1
	Thrashing (moving wildly)	2
Agitation	Rigid (stiff)	2
	Asleep or clam	0
	Can be comforted to lessen the agitation (mild)	1
Complains of pain	Can not be comforted (hysterical)	2
	Asleep	0
	States no pain	0
	Can not localize	1
	Localizes pain	2

Table4: Oral analgesia

	Time of First analgesia (hrs.)	No. of doses in 24 hrs.
Group A	183+16.5	6.00+ 00
Group B	350+20.5	1.63+.597
Group C	594+148.12	1.40+.502

Table 5: Modified Alderte Recovery Scoring System

Parameter	Finding	Points
Activity: able to move, voluntarily or command	Four extremities	2
	Two extremities	1
	No extremities	0
Respiration	Able to breathe deeply and cough freely	2
	Dyspnoea, shallow or limited breathing	1
	Apnoea	0
Circulation	Blood Pressure within 20 mm Hg of preoperative level	2
	Blood Pressure within 20-50 mm Hg of preoperative level	1
	Blood Pressure + 50 mm Hg of preoperative level	0
Consciousness	Fully awake	2
	Arousable on calling	1
	Unresponsive	0
Oxygen Saturation	Saturation >92%	2
	Needs oxygen to maintain saturation >90%	1
	Saturation <90% with oxygen	0

Nine or more points are required for recovery to be confirmed

DISCUSSION

Lower abdominal and lower limb surgeries in children are associated with considerable postoperative Pain. In this study, the authors evaluated the efficacy of fentanyl and ketamine as adjuvants to bupivacaine in caudal analgesia in comparison to that of bupivacaine alone in children undergoing genitourinary and lower limb surgeries. Many previous researchers⁽¹⁹⁻²¹⁾ have observed that caudal block with local anesthetics alone provides adequate intra-operative analgesia in only 57-84% of children, whereas 93-100% of children had adequate analgesia when caudal block with fentanyl 1 mcg/kg was used. The efficacy of caudal block for intra-operative analgesia can be affected by a number of factors. Such as the volume and concentration of local anesthetic used, the surgical procedure, criteria used to define adequate analgesia and the depth of general anesthesia⁽²²⁾. Locatelli et al⁽⁷⁾ and Breschan et al⁽⁸⁾ concluded from their study that bupivacaine and levobupivacaine are equally potent and had longer analgesic effect. Hence, 0.25% bupivacaine was chosen for the present study. In contrast to the definition of adequate analgesia adopted in the present study, Constant et al⁽¹⁹⁾ and Martindale and co-workers⁽²³⁾ used an increase by 15% in the hemodynamic parameters as criteria for adequate caudal block. Constant and colleagues⁽¹⁹⁾ maintained anesthesia with 0.6 MAC of isoflurane, whereas in the present study anesthesia was maintained using 1% sevoflurane and 67% nitrous oxide at the time of surgical incision. The deeper plane of anesthesia in our study may have offered stable hemodynamic parameters.

No statistical differences were observed in all the groups with regard to intra-operative. Hemodynamic changes except for a significant decrease

in heart rate in group B which received fentanyl. The decrease in heart rate was not associated with hypotension and did not require any intervention. (Which may be attributed to the systemic effects of fentanyl).

In the present study, the mean duration of post-operative analgesia from the time of extubation to the first administration of rescue analgesic in group A was 183±16.57 minutes. This was in contrast to a study which observed 5 hours of post-operative analgesia using 1ml/kg of bupivacaine for caudal block.⁽²⁴⁾ The mean duration of post-operative analgesia in group B (350±20.5 minutes) was consistent with previous studies.⁽²⁴⁻²⁵⁾ By reducing the dose of Ketamine from 1 mg/kg to 0.5 mg/kg the incidence of side effect were reduced⁽²⁴⁾ by 5% but the duration of analgesia obtained in group C was significantly longer than that in groups A and B. Our findings are consistent with those of Cook and colleagues⁽²⁴⁾ who demonstrated that the addition of Ketamine 0.5 mg/kg to bupivacaine 0.25%, 1 ml/kg provided longer duration of analgesia than bupivacaine alone and behavioral side effects attributable to ketamine were not observed. We chose to monitor patients for a period of 24 hours post-operatively. This is in contrast to a few other studies^(26,27) where there was only a six-hour period of observation. Post-operatively and the rest of the assessment was done by parents which could introduce some inconsistencies, as parents differ in the way they perceive their children to be in pain and the threshold for administering rescue medications varies between parents.⁽²⁸⁾

CONCLUSION

Addition of fentanyl 1 mcg/kg or ketamine 0.5 mg/kg to caudal bupivacaine prolongs postoperative analgesia in children undergoing genitourinary and lower limb surgery. Caudal fentanyl with bupivacaine does not increase the incidence of post-operative respiratory depression. Post-operative pain relief is prolonged significantly with ketamine as compared to fentanyl.

REFERENCES

- De Beer DA, Thomas ML. Caudal additives in children-- solutions or problems? *Br J Anaesth.* 2003;90(4):487-98.
- Lloyd-Thomas AR. Pain management in paediatric patients. *Br J Anaesth.* 1990;64(1):85-104.
- Sanders JC. Paediatric regional anaesthesia, a survey of practice in the United Kingdom. *Br J Anaesth.* 2002;89(5):707-10.
- Whelton A. Nephrotoxicity of nonsteroidal anti-inflammatory drugs: physiologic foundations and clinical implications. *Am J Med.* 1999;106(5):13S-24S.
- Maxwell LG, Kaufmann SC, Bitzer S, Jackson Jr EV, et al. The effects of a small-dose naloxone infusion on opioid-induced side effects and analgesia in children and adolescents treated with intravenous patient-controlled analgesia: a double-blind, prospective, randomized, controlled study. *Anesthesia & Analgesia.* 2005;100(4):953-958.
- Frawley GP, Downie S, Huang GH. Levobupivacaine caudal anesthesia in children: a randomized double-blind comparison with bupivacaine. *Paediatr Anaesth.* 2006; 16:754-60
- Locatelli B, Ingelmo P, Sonzogni V, Zanella A, Gatti V, Spotti A, et al. Randomized, double-blind, phase III, controlled trial comparing levobupivacaine 0.25%, ropivacaine 0.25% and bupivacaine 0.25% by the caudal route in children. *Br J Anaesth.* 2005;94:366-71.
- Breschan C, Jost R, Krumpolz R, Schaumberger F, Stettner H, Marhofer P, et al. A prospective study comparing the analgesic efficacy of levobupivacaine, ropivacaine and bupivacaine in pediatric patients undergoing caudal blockade. *Paediatr Anaesth.* 2005; 15:301-6.
- Kumar P, Rudra A, Pan AK, Acharya A. Caudal additives in pediatrics: A comparison among Midazolam, ketamine and neostigmine co administered with bupivacaine. *Anesth Analg* 2005;101:69-73
- Engelman E, Marsala C. Bayesian enhanced meta-analysis of post-operative analgesic efficacy of additives for caudal analgesia in children. *Acta Anaesthesiol Scand.* 2012;56:817-32
- Demiraran Y, Kocaman B, Akman Y. A comparison of the postoperative analgesic efficacy of single-dose epidural tramadol versus morphine in children. *Br J Anaesth.* 2005 ;95:510-3.
- Gunes Y, Gunduz M, Unlugenc H, Ozalevi M, Ozcengiz D. Comparison of caudal vs intravenous tramadol administered either preoperatively or postoperatively for pain relief in boys. *Paediatr Anaesth.* 2004; 14:324-8
- Senel AC, Akyol A, Dohman D, Solak M. Caudal bupivacaine-tramadol combination for postoperative analgesia in pediatric herniorrhaphy. *Acta Anaesthesiol Scand.* 2001;45:786-9
- Yildiz T, Ozdamar D, Bagus F, Solak M, Toker K. Levobupivacaine-tramadol combination for caudal block in children: a randomized, double-blinded, prospective study. *Paediatr Anaesth.* 2010;20:524-9.
- Naguib M, Sharif AMY, Seraj M, El Gamma! M, Dawlatly AA. Ketamine for caudal analgesia in children: comparison with caudal bupivacaine. *Br J Anaesth.* 1991 ;67(5):559-64.
- Ahuja S, Yadav S, Joshi N, Chaudhary S, Madhu SV. Efficacy of caudal fentanyl and ketamine on post-operative pain and neuroendocrine stress response in children undergoing infraumbilical and perineal surgery: A pilot study. *Journal of anaesthesiology, clinical pharmacology.* 2015 ;31(1): 104.
- Bans S, Karakaya D, Kelsaka E, Guldogus F, Ariturk E, Tür A. Comparison of fentanyl bupivacaine or midazolam—bupivacaine mixtures with plain bupivacaine for caudal anaesthesia in children. *Pediatric Anesthesia.* 2003;13(2):126-131.
- Aldrete JA. The post anaesthesia recovery score revisited. *J. clin Anaesth* 1995 feb; 7(1) : 89-91
- Hannallah R, Broadman L. Comparison of caudal and ilioinguinal/iliohypogastric nerve blocks for control of post-orchiopey pain in pediatric ambulatory surgery. *Anesthesiol.* 1987; 66: 832-834.
- Constant I, Gall O, Gouyet L, Chauvin M, Murat I. Addition of clonidine or fentanyl to local anaesthetics prolongs the duration of surgical analgesia after single shot caudal block in children. *Br J Anaesth.* 1998 Mar;80(3):294-8.
- Semple D, Findlow D, Aldridge LM, Doyle E. The optimal dose of ketamine for caudal epidural blockade in children. *Anaesthesia.* 1996;51(12):1170-72.
- Ahuja S, Yadav S, Joshi N, Chaudhary S, Madhu SV. Efficacy of caudal fentanyl and ketamine on post-operative pain and neuroendocrine stress response in children undergoing infraumbilical and perineal surgery: A pilot study. *Journal of anaesthesiology, clinical pharmacology.* 2015 ;31(1): 104.
- Senel AC, Akyol A, Dohman D, Solak M. Caudal bupivacaine-tramadol combination for postoperative analgesia in pediatric herniorrhaphy. *Acta anaesthesiologica scandinavica.* 2001 ;45(6):786-89.
- Martindale SJ, Dix P, Stoddart PA. Double-blind randomized controlled trial of caudal versus intravenous S (+)-ketamine for supplementation of caudal analgesia in children. *Br J Anaesth.* 2004;92(3):344-47.
- Campbell FA, Yentis SM, Fear DW, Bissonnette B. Analgesic efficacy and safety of a caudal bupivacaine-fentanyl mixture in children. *Can J Anaesth.* 1992;39(7):661-64.
- Cook B, Grubb DJ, Aldridge LA, Doyle E. Comparison of the effects of adrenaline, clonidine and ketamine on the duration of caudal analgesia produced by bupivacaine in children. *Br J Anaesth.* 1995;75(6):698-701.
- Kaul A, Pant D, Sood J. Caudal clonidine in day-care paediatric surgery. *Ind j Anaesth.* 2009;53(4):450.
- Klimscha W, Chiari A, Michalek-Sauberer A, Wildling E, Lerche A, Lorber C et al. The efficacy and safety of a clonidine/bupivacaine combination in caudal blockade for pediatric hernia repair. *Anesthesia & Analgesia.* 1998;86(1):54-61.
- Parameswari A, Dhev AM, Vakamudi M. Efficacy of clonidine as an adjuvant to bupivacaine for caudal analgesia in children undergoing sub-umbilical surgery. *Ind j Anaesth.* 2010;54(5):458.