

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Clinical Anaesthesia

Journal homepage: [www.ijca.in](http://www.ijca.in)

## Review Article

# A systematic review on efficacy of dexmedetomidine on emergence agitation after nasal surgeries

Ritu Medatwal<sup>1\*</sup>, Anil Garg<sup>1</sup>

<sup>1</sup>Dept. of Anaesthesiology, Amaltas Institute of Medical Science, Bangar, Madhya Pradesh, India



## ARTICLE INFO

### Article history:

Received 20-02-2023

Accepted 22-03-2023

Available online 26-03-2024

### Keywords:

Nasal surgery

Agitation

Emergence

Dexmedetomidine

## ABSTRACT

Dexmedetomidine has sympatholytic, soothing, pain relieving vasoconstrictive and, sedative impacts, which might assist with forestalling hypotension. This study intended to play out an orderly survey of the writing and explore the viability of dexmedetomidine on perioperative dizziness following nasal medical procedure. We looked the electronic Embase, Cochrane, MEDLINE, and PubMed data sets. Moreover, each significant article's and book's reference index was totally looked. Included were the expressions "Dexmedetomidine" [MeSH] "development disturbance," [MeSH], "nasal medical procedure" [MeSH] and general anaesthesia" [MeSH]. This survey shows that the fundamental organization of dexmedetomidine can diminish intraoperative blood misfortune, careful time, postoperative torment and the rate of development unsettling.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Rise agitation, otherwise called development ridiculousness, or post-sedative fervor, is a clinical peculiarity happening in the quick postoperative period following general anaesthesia.<sup>1</sup> However it is most regularly found in kids & the elderly<sup>2,3</sup> can influence all age gatherings. Development is the momentary cycle from obviousness to full wakefulness<sup>4</sup>, & this progress from general sedation is normally smooth & uneventful.<sup>5</sup> Rise fomentation is typically seen during the initial 15-30 minutes following awakening<sup>3,4</sup> & has a variable clinical show like fretfulness, peevishness, whipping, unsettling, hyperactivity, crying, groaning, confusion & uncooperative behaviour.<sup>3-6</sup> However rise disturbance is very much perceived clinically, it should be better perceived., Since there is trouble separating among tumult & torment, & vulnerability about its clinical importance needs, more

exact definitions are dependable with substantial evaluation tools.<sup>2</sup> The rate reported in the pediatric populace is 12-13%<sup>3</sup> & around 10-15% in the geriatric age group.<sup>7,8</sup> There are a few gamble factors related with EA like the presence of an endotracheal tube or urinary catheter, premedication with benzodiazepine, inhalational sedatives,<sup>3,6</sup> bosom, abdominal & otolaryngologic surgeries.<sup>7,8</sup> Development unsettling can prompt entanglements like self-extubation & expulsion of catheters, yearning pneumonia dying, patients might encounter hypertension & tachycardia & protracts the clinic stay.<sup>2,9,10</sup> It can increment medical care costs as it requires expanded staff, drugs, & delayed PACU stay.<sup>3,9</sup> Among the different medical procedures, otolaryngologic methodology are related with a higher occurrence of agitation.<sup>7,8,11</sup> It is seen that these patients might have a sensation of suffocation<sup>9</sup> during rise because of the presence of nasal pressing. Different methodology have been utilized to forestall or treat EA. Recognizing & killing the encouraging elements is the most fundamental among them. Pharmacological specialists

\* Corresponding author.

E-mail address: [rmedatwal@gmail.com](mailto:rmedatwal@gmail.com) (R. Medatwal).

like propofol, opioids,<sup>12–14</sup> clonidine,<sup>15</sup> midazolam,<sup>16</sup> & dexmedetomidine<sup>14,17,18</sup> were utilized prophylactically to decrease the rate of rise fomentation. Dexmedetomidine is a specific alpha-2 adrenergic agonist with pain relieving, narcotic & sympatholytic properties. It doesn't follow up on GABA receptors & consequently doesn't create respiratory misery. Also, it has the interesting property of fast beginning with brief length helpful sedation. It is likewise antisecretory & anxiolytic.<sup>19</sup> It has regularly been utilized to diminish EA in children<sup>20–23</sup> & for weaning from the ventilator in escalated care unit.<sup>24</sup> It decreases torment power, narcotic utilization, & stress reaction & hence further develops recuperation after surgery.<sup>17,25–27</sup> The writing concerning the impacts of dexmedetomidine on lessening rise disturbance in grown-ups is restricted & less in number.

## 2. Materials and Methods

This study was performed according to the Preferred Reporting Items for Systematic Review. All studies regarding the effect of dexmedetomidine on emergence agitation published in the English language peer-reviewed scientific journals from April 2013 to September 2021, studies in which the outcome was defined in terms of efficacy of dexmedetomidine, all the articles published till 31 September 2021 & full articles in English were included in the study for sample size selection.

All the case reports, case series, cross-sectional studies, reviews, abstracts & articles with incomplete data & patients with lesions were excluded. All the articles were published in any other language except English. The references of selected articles were also analyzed for additional studies & any study that did not meet the inclusion criteria.

### 2.1. Search strategy

We looked the electronic Embase, Cochrane, MEDLINE, and PubMed data sets. Furthermore, each pertinent article's and book's reference index was totally looked. The appropriate papers were picked by two analysts independently founded on the rules. The two analysts discussed any distinctions until they concurred.

### 2.2. Selection

The determination of the examinations were acted in . To start with, all book titles were analyzed, and reasonable examinations were picked per the consideration and rejection standards. Abstracts for every one of the picked titles were gained, inspected, and significant digests were picked in view of the measures. At long last, the conclusive assortment of articles was gotten while remembering the determination standards after full-text variants of the relative multitude of digests that had been picked had been

acquired and analyzed, which is displayed in Table 1.

**Table 1:** Selection of articles

Initial search	180
Duplicates & irrelevant	71
Case reports & series	15
Reviews	62
Abstract	13
Language other than English	13

Five articles were selected for the study.

### 2.3. Quality assessment

The adequacy of the examinations was assessed utilizing the Cochrane Coordinated effort Apparatus for assessing the Gamble of Predisposition in RCTs. Instances of choice inclination incorporate arbitrary grouping age and assignment covering, execution predisposition, whittling down predisposition, announcing predisposition, and some other bias found. Each chose concentrate on got a high, low, or uncertain gamble evaluation. Three onlookers freely surveyed the situation, and any conflicts were talked about.

## 3. Results

Initial searches yielded 100 articles. Five studies were considered for analysis out of a total of 100 articles that were found in the database after duplicates were removed & publications that did not meet eligibility requirements were eliminated. The PRISMA flowchart for the inclusion of studies is shown in Figure 1.

### 3.1. Synthesis of results

Account amalgamation has been accommodated the discoveries gotten from the examinations. The information extricated has been introduced in the even structure as referenced underneath. Table 2 showed the included examinations and Table 3 showed the positive discoveries of the review.

**Table 2:** Included studies


Author	Type of Study	Country
S Y Kim et al, 2013 <sup>10</sup>	RCT	Seoul, Korea
R Polat et al, 2015 <sup>28</sup>	RCT	Ankara, Turkey
Hina Khurshid et al, 2015 <sup>29</sup>	RCT	Srinagar, India
Akaansha Garg et al, 2018 <sup>30</sup>	RCT	Jodhpur, India
Ramesh Kumar N et al, 2020 <sup>31</sup>	RCT	Karnataka, India

**Table 3:** Positive findings of the included studies

Author	Sample Size	Result obtained	Conclusion
S Y Kim et al, 2013 <sup>10</sup>	100	The occurrence of disturbance and mean blood vessel pressure and pulse were lesser in bunch with was lower in dexmedetomidine .Time to extubation, bispectral file, and respiratory rate at extubation were comparable between the gatherings.	Intraoperative imbuement of dexmedetomidine gave smooth and stable development. It likewise worked on nature of recuperation (Qor=40) after nasal medical procedure.
R Polat et al, 2015 <sup>28</sup>	90	The occurrence of EA was altogether higher in bunch with dexmedetomidineas (46.7%) contrasted with bunch remifentanil (3.3 and 20%, separately; p < 0.001). Lingering sedation in the post-sedation care unit (PACU) was comparative in all gatherings (p = 0.947). The occurrence of queasiness or regurgitating was altogether lower in bunch dexmedetomidineas	Anesthetic maintenance with either remifentanil or dexmedetomidine implantation until extubation gave a more smooth and hemodynamically stable development.
Hina Khurshid et al, 2015 <sup>29</sup>	100	Rate of rise disturbance was lower in Gathering dexmedetomidin (26%) . Bunch dexmedetomidin showed more steady hemodynamics than Gathering II.	The use of intraoperatively infused dexmedetomidine as resulted in smooth emergence with stable hemodynamics
Akaansha Garg et al, 2018 <sup>30</sup>	72	Dexmedetomidine mixture essentially diminished the rate of EA (Gathering C 52.8%; Gathering D 5.6%) by 89.5% (P = 0.00001). The end-flowing desflurane fixation was lower, and there was a 28.87% decrease in desflurane necessity in bunch D contrasted with bunch C (P < 0.001).	Dexmedetomidine significantly reduced the incidence of EA & requirement of desflurane in patients undergoing nasal surgery.
Ramesh Kumar N et al, 2020 <sup>31</sup>	100	All segment information were equivalent between the two gatherings. Disturbance score $\geq 5$ was seen in 4% and 58% and was genuinely huge (p<0.5). Time to verbal reaction was more in the dexmedetomidine bunch (6.74 $\pm$ 2.13 min v/s 5.78 $\pm$ 1.91 min)	Dexmedetomidine as an infusion reduces emergence agitation in adults undergoing nasal surgery under desflurane general anaesthesia

**Table 4:** Risk of bias in the included studies

Authors name	Selection Bias. Random sequence generation	Allocation Concealment	Reporting bias	Others	Performance bias. Blinding participants & personnel	Blinding Outcome	Attrition bias
S Y Kim et al, 2013 <sup>10</sup>	Low risk	Low risk	Low risk	Low risk	Low risk	Unclear	High risk
R Polat et al, 2015 <sup>28</sup>	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Hina Khurshid et al, 2015 <sup>29</sup>	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk
Akaansha Garg et al, 2018 <sup>30</sup>	Low risk	Unclear	Low risk	Low risk	Low risk	Unclear	Low risk
Ramesh Kumar N et al, 2020 <sup>31</sup>	Low risk	Low risk	Low risk	Low risk	Unclear	Unclear	Low risk



**Figure 1:** PRISMA flowchart

### 3.2. Risk of bias assessment

The risk of inclination was assessed utilizing the Cochrane Chance of Predisposition Appraisal Instrument. Every component from one of five spaces is given an inclination score (high, low, or hazy) (determination, execution, whittling down, revealing, & other). Part I of the structure (Quality appraisal) assesses the gamble of choice, revealing, & other predisposition while to some degree II, hazard of execution, discovery, & weakening predisposition are evaluated., Risk of inclination was characterized for every

judgment as "high," "low," or "muddled" involving the directions at the lower part of the survey as referenced in Table 4.

### 4. Discussion

Development agitation is a very much depicted clinical, complex peculiarity of multifactorial etiology from rise till general sedation. Albeit clinical examinations exploring EA keep on being distributed, this field faces impediments because of the requirement for highest

quality level appraisal instrument. In this manner looking at results across changed examinations stays risky & refreshed clinical administration apparatuses are required. Different pharmacological specialists have been utilized prophylactically to lessen EA following general sedation with variable achievement rates.<sup>10,12–16,18,21</sup> The job of benzodiazepine premedication on EA is questionable. A few examinations revealed that the utilization of benzodiazepine premedication inclines toward EA, while others have viewed it as defensive. Preoperative uneasiness is likewise a realized gamble factor for EA.<sup>9</sup> Benzodiazepines are most ordinarily used to preoperatively decrease tension. Given the above reasons, we involved midazolam for premedication in all patients per the institutional practice. Dexmedetomidine is one among the different pharmacological specialists utilized prophylactically for EA.<sup>10,18,21</sup> It has been utilized as a solitary bolus infusion, consistent mixture or bolus followed by persistent implantation in different portions to forestall EA with variable success.<sup>10,18,21</sup> The suitable portion where it is viable in forestalling EA is yet to be distributed in the writing. No "best quality level" appraisal apparatus is accessible to assess EA. The different scales used to assess tumult are; the Ricker disturbance sedation scale, Richmond sedation unsettling scale, the engine action evaluation scale & the altered New Sheffield sedation scale. Studies have shown that the Ricker & Richmond scales have fantastic between rater unwavering quality. This large number of scales were made for grown-ups in the ICU, & none has been laid out exclusively for use in PACU. Nonetheless, we feel patients owned up to the PACU are tantamount to those confessed to the ICU, & these scales are not difficult to utilize & give an exact assessment of fomentation. Rise tumult is boundless after ENT surgeries,<sup>7,8,11</sup> & this higher frequency is credited to a feeling of suffocation because of nasal packing.<sup>11</sup> The occurrence of EA for nasal medical procedures detailed in the writing changes from 22.2% to 55.4%.<sup>7</sup> Dexmedetomidine is known to diminish the perioperative necessity of narcotics & has a narcotic saving activity.

## 5. Conclusion

Intraoperative dexmedetomidine infusion lessens the rate of EA in the quick postoperative period in grown-up patients going through nasal medical procedure under desflurane sedation. In any case, it could prompt postponed recuperation as far as delayed chance to extubation, time to accomplish BIS worth of 90, time to patient reaction on verbal order & time to release from the postanesthesia care unit. The fundamental organization of dexmedetomidine can diminish careful time, intraoperative blood misfortune, & portions of intraoperative breathed in sedative gas & fentanyl contrasted & a fake treatment. It can likewise diminish postoperative torment & the frequency of crisis unsettling. Because of less examinations, further clinical

preliminaries are expected to affirm these outcomes.

## 6. Source of Funding

None.

## 7. Conflict of Interest

None.

## References

- O'Brien D. Acute postoperative delirium: definition, incidence, recognition & interventions. *J Perianesth Nurs*. 2002;17(6):384–92.
- Scott GM, Gold JJ. Emergence delirium: a re-emerging interest. *Semin Anesth Perioper Med Pain*. 2006;25(3):100–4.
- Hudek K. Emergence delirium: a nursing perspective. *AORN J*. 2009;89(3):509–16.
- Xara D, Silva A, Mendonca J, Abelha F. Inadequate emergence after anaesthesia: emergence delirium & hypoactive emergence in the Postanaesthesia Care Unit. *J Clin Anesth*. 2013;25(6):439–46.
- Radtke FM, Frank M, Hagemann L, Seeling M, Wernecke KD, Spies CD. Risk factors for inadequate emergence after anaesthesia: emergence delirium & hypoactive emergence. *Minerva Anesthesiol*. 2010;76(6):394–403.
- Burns SM. Delirium during emergence from anaesthesia: a case study. *Crit Care Nurse*. 2003;23(1):66–9.
- Yu D, Chai W, Sun X, Yao L. Emergence agitation in adults: risk factors in 2,000 patients. *Can J Anaesth*. 2010;57(9):843–8.
- Lewis TV, Malviya S, Tait AR. A prospective cohort study of emergence agitation in the pediatric postanesthetic care unit. *Anesth Analg*. 2003;96(6):1625–30.
- Lepouse C, Lautner CA, Liu L, Gomis P, Leon A. Emergence delirium in adults in the post-anaesthesia care unit. *Br J Anaesth*. 2006;96(6):747–53.
- Kim SY, Kim JM, Lee JH, Song BM, Koo BN. Efficacy of intraoperative dexmedetomidine infusion on emergence agitation & quality of recovery after nasal surgery. *Br J Anaesth*. 2013;111(2):222–8.
- Eckenhoff JE, Kneale DH, Dripps RD. The incidence & etiology of postanesthetic excitement. A clinical survey. *Anesthesiology*. 1961;22:667–73.
- Cohen IT, Hannallah RS, Hummer KA. The incidence of emergence agitation associated with desflurane anesthesia in children is reduced by fentanyl. *Anesth Analg*. 2001;93(1):88–91.
- Galinkin JL, Fazi LM, Cuy RM, Chiavacci RM, Kurth CD, Shah UK, et al. Use of intranasal fentanyl in children undergoing myringotomy & tube placement during halothane & sevoflurane anesthesia. *Anesthesiology*. 2000;93(6):1378–83.
- Finkel JC, Cohen IT, Hannallah RS, Patel KM, Kim MS, Hummer KA, et al. The effect of intranasal fentanyl on the emergence characteristics after sevoflurane anesthesia in children undergoing surgery for bilateral myringotomy tube placement. *Anesth Analg*. 2001;92(5):1164–8.
- Kulka PJ, Bresslem M, Tryba M. Clonidine prevents sevoflurane-induced agitation in children. *Anesth Analg*. 2001;93(2):335–8.
- Ko YP, Huang CJ, Hung YC, Su NY, Tsai PS, Chen CC, et al. Premedication with lowdose oral midazolam reduces the incidence & severity of emergence agitation in pediatric patients following sevoflurane anesthesia. *Acta Anaesthesiol Sin*. 2001;39(4):169–77.
- Gurbet A, Mongol EB, Turker G, Ugun F, Kaya FN, Ozcan B. Intraoperative infusion of dexmedetomidine reduces perioperative analgesic requirements. *Can J Anaesth*. 2006;53(7):646–52.
- Ibacahe ME, Munoz HR, Brandes V, Morales AL. Single-dose dexmedetomidine reduces agitation sevoflurane anaesthesia in children. *Anesth Analg*. 2004;98(1):60–3.
- Gerlach T, Dasta JF. Dexmedetomidine: an updated review. *Ann Pharmacother*. 2007;41(2):245–52.

20. Patel A, Davidson M, Tran MCJ, Quraishi H, Schoenberg C, Sant M, et al. Dexmedetomidine infusion for analgesia & prevention of emergence agitation in children with obstructive sleep apnea syndrome undergoing tonsillectomy & adenoidectomy. *Anesth Analg*. 2010;111(4):1004–10.
21. Shukry M, Clyde MC, Kalarickal PL, Ramadhyani U. Does dexmedetomidine prevent emergence agitation after sevoflurane-based anaesthesia? *Paediatr Anaesth*. 2005;15(12):1098–104.
22. Guler G, Akin A, Tosun Z, Ors S, Esmaglu A, Boyaci A. Single dose dexmedetomidine reduces agitation & provides smooth extubation after pediatric adenotonsillectomy. *Paediatr Anaesth*. 2005;15(9):762–6.
23. Erdil R, Demirebilek S, Begec Z, Ozturk E, Ulger MH, Ersoy MO. The effects of dexmedetomidine & fentanyl on emergence characteristics after adenoidectomy in children. *Anaesth Intensive Care*. 2009;37(4):571–6.
24. Shehabi Y, Nakae H, Hammond N, Bass F, Nicholson L, Chen J. The effect of dexmedetomidine on agitation during weaning of mechanical ventilation in critically ill patients. *Anaesth Intensive Care*. 2010;38(1):82–90.
25. Tufanogullari B, White PF, Peixoto MP, Kianpour D, Lacour T, Griffin J, et al. Dexmedetomidine infusion during laparoscopic bariatric surgery: The effect on recovery outcome variables. *Anesth Analg*. 2008;106(6):1741–8.
26. Bekker A, Haile M, Kline R. The effect of intraoperative infusion of dexmedetomidine on quality of recovery after major spinal surgery. *J Neurosurg Anesthesiol*. 2013;25(1):16–24.
27. Blandszun G, Lysakowski C, Elia N, Tramèr MR. Effect of perioperative systemic  $\alpha_2$  agonists on perioperative morphine consumption & pain intensity: systemic review & meta-analysis of randomized controlled trials. *Anesthesiology*. 2012;116(6):1312–22.
28. Polat R, Peker K, Baran I, Aydın GB, Gülöksüz Ç, Dönmez A. Comparison between dexmedetomidine and remifentanyl infusion in emergence agitation during recovery after nasal surgery: A randomized double-blind trial. *Anaesthesist*. 2015;64(10):740–6.
29. Khurshid H, Muneer K, Malla MS. Effect of Dexmedetomidine on Emergence Agitation after Nasal Surgeries. *Indian J Clin Anaesth*. 2015;2(3):126–30.
30. Garg A, Kamal M, Mohammed S, Singariya G, Chouhan DS, Biyani G. Efficacy of dexmedetomidine for prevention of emergence agitation in patients posted for nasal surgery under desflurane anaesthesia: A prospective double-blinded randomised controlled trial. *Indian J Anaesth*. 2018;62(7):524–30.
31. Kumar RN. Dexmedetomidine infusion for prevention of emergence agitation after nasal surgery. *J Evol Med Dent Sci*. 2020;9(11):844–50.

### Author biography

**Ritu Medatwal**, Assistant Professor

**Anil Garg**, Associate Professor

**Cite this article:** Medatwal R, Garg A. A systematic review on efficacy of dexmedetomidine on emergence agitation after nasal surgeries. *Indian J Clin Anaesth* 2024;11(1):78-83.